

## **SITE SPECIFIC BINDING SYSTEM, IMAGING COMPOSITIONS AND METHODS**

### **CROSS-REFERENCE TO RELATED APPLICATIONS**

**[0001]** This application is a divisional of application Serial No. 09/404,963 which is a continuation-in-part of application Serial No. 09/404,963, filed September 24, 1999 which is a continuation-in-part of application Serial No. 09/189,118, filed November 9, 1998 which is a continuation of application Serial No. 08/854,308, filed May 12, 1997 which is a divisional application of Serial No. 08/488,743 filed June 8, 1995, now U.S. patent no. 5,690,907 dated November 25, 1997.

### **BACKGROUND OF THE INVENTION**

**[0002]** This invention relates to a novel site specific binding system and novel compositions, and more particularly, to such a system and compositions which are useful in improved methods for ultrasonic imaging, drug or chemotherapeutic agent delivery, and diagnostic assays and detection systems.

**[0003]** Heretofore, with respect to ultrasonic imaging, although ultrasonic contrast agents based upon "bubble" technology have been demonstrated to develop an acoustic impedance mismatch by virtue of gas encapsulated either in protein (Feinstein et al., J. Am. Coll. Cardiol. 1990; 16:316-324 and Keller et al., J. Am. Soc. Echo. 1989; 2:48-52), polysaccharide (Corday et al., J. Am. Coll. Cardiol. 1984; 3:978-85) biodegradable polymers (Schneider et al., Invest. Radiol., 1993; 27:134-139 and Bichon et al., European Patent Application No. 8908103 67.4: 1990) or lipids (D'Arrigo et al., J. Neuromag., 1991; 1: 134139; Simon et al., Invest. Radiol., 1992; 27:29-34; and Unger et al., Radiology 1992; 195:453-456), no experimental evidence of site-specific targeting of an acoustic contrast or imaging agent with resultant changes in the acoustic properties of the targeted tissue, surface or support are known. This lack of results has occurred despite numerous methods described in the literature for modifying such agents for targeting purposes, and the failure of past targeting approaches may be due to the chemical nature of the agents, production process limitations or particle instabilities.

**[0004]** Nongaseous acoustic contrast agents have been described including lipid emulsions(Fink et al., Ultrason. Imaging, 1985 7:191-197) liposomes

(Lanza et al., J. Am. Coll. Cardiol., 1992 (abstract); 19 (3 Suppl A) 114A), and perfluorocarbon emulsions (Mattrey et al., Radiology 1982; 145: 759-762 and Mattrey et al., Ultrasound Med. 1983; 2:173-176). As with the contrast agents discussed above, no demonstration of site targeted emulsion or liposome has been reported. Again, such failure may reflect instability of the particles, process incompatibilities or the chemical nature of the contrast agent. Lipid emulsions were evaluated by Fink et al. supra and did not exhibit adequate echogenicity in studies examining hepatic imaging. A unique chemical formulation of liposomes described by Lanza et al. supra was suggested to have the potential to be a targetable ultrasonic contrast but such has not been demonstrated to date. Perfluorocarbon emulsions, Perflubron (perfluorooctylbromide, P 100) and Flusol (perfluorodecalin and perfluorotripropylamine, F20) have been used as ultrasonic contrast agents and have been reported to accumulate in liver, spleen and tumors secondary to phagocytic uptake of emulsion particles at these sites (Mattrey et al. 1983, supra). These perfluorocarbon emulsions have also been noted to enhance Doppler signals and opacify lumens. Fluorocarbons and fluorocarbon emulsions for use as contrast agents are disclosed in U.S. Patent Nos. 4,927,623, 5,077,036, 4,838,274, 5,068,098, 5,114,703, 5,362,477, 5,362,478, 5,171,755, 5,304,325, 5,350,571 and 5,403,575. However, no demonstration of perfluorocarbon emulsions as a ligand targeted acoustic contrast system has been reported.

**[0005]** Previous descriptions of tissue or organ targeting in biomedical ultrasonics has referred to the collection of acoustically reflective particles within or around structural tissue abnormalities. Localized acoustic enhancement of tissue pathologies (e.g. malignancies) has not been ligand-directed but rather has depended upon differential dynamic rates of particle uptake and/or clearance between normal and malignant tissues. Such contrast agents have included aqueous solutions (Ophir et al., Ultrason. Imaging 1979, 1:265-279; Ophir et al., Ultrasound Med. Biol. 1989, 15:319-333; and Tyler et al., Ultrason. Imaging, 3:323-329), emulsions (Fink et al. Ultrason. Imaging, 1985, 7:191-197), and suspensions (Mattrey et al. 1982 supra and Mattrey et al., Radiology, 1987, 163:339-343). Although the possibility of ligand-directed ultrasonic contrast targeting with acoustically reflective liposomes has been suggested, no successful applications of this concept have been reported (Lanza et al. 1992, supra and Valentini et al., J. Am. Coll. Cardiol., 1995, 25:16A). Previous approaches to targeting in vivo of particles have involved direct conjugation of a

ligand (e.g. monoclonal antibody) to a vesicle by a variety of methods (see, for example, Torchlin et al., Biochem. Biophys. Res. Commun. 1978, 85:983-990; Endoh et al., J. Immunol. Methods, 1981, 44:7985; Hashimoto et al., J. Immunol. Methods, 1983, 62:155-162 and Martin et al., Biochemistry, 1981, 20:4229-4238).

**[0006]** There remains a need for new and improved methodologies for ligand-based binding systems which can be adapted as an ultrasonic contrast system permitting detection of molecular moieties such as peptides, carbohydrates or nucleic acids and whose uses can range from ultrasound-based ELISA-like laboratory diagnostic assays in liquid and solid phase systems and in cell cultures; electrophoretic, chromatographic and hybridization detection systems to the detection of thrombi, infections, cancers and infarctions in patients with the use of conventional ultrasonic imaging methods.

### **SUMMARY OF THE INVENTION**

**[0007]** Among the several objects of the invention may be noted the provision of a novel method for ligand-based binding of lipid encapsulated particles to molecular epitopes on a surface in vivo or in vitro, the provision of such a method in which the ligand is conjugated to the lipid encapsulated particles through an avidin-biotin interaction and the resulting conjugate is bound to molecular epitopes on a surface; the provision of such a method which is useful for enhancing the acoustic reflectivity of a biological surface for ultrasonic imaging; the provision of a method of this type wherein the conjugate formed is effective for imaging by x-ray, ultrasound, magnetic resonance or positron emission tomography; the provision of compositions for use in ultrasonic imaging of a biological surface and for enhancing the acoustic reflectivity of such a surface; the provision of ultrasonic contrast agents which become highly reflective when bound to the desired site or biological surface through the ligand-based binding system of the invention; and the provision of such methods and compositions which are capable of targeting and altering the echogenic properties of a tissue surface for improved and specific identification of pathological processes. Other objects will be in part apparent and in part pointed out hereinafter.

**[0008]** Briefly, in its broadest embodiment, the present invention is directed to a method for ligand-based binding of lipid encapsulated particles to molecular epitopes on a surface in vivo or in vitro which comprises sequentially administering (a) a site-specific ligand activated with a biotin activating agent; (b) an

avidin activating agent; and (c) lipid encapsulated particles activated with a biotin activating agent, whereby the ligand is conjugated to the particles through an avidin-biotin interaction and the resulting conjugate is bound to the molecular epitopes on such surface. The conjugate is effective for imaging by x-ray, ultrasound, magnetic resonance or positron emission tomography. In a more specific embodiment, the invention is directed to a method for enhancing the acoustic reflectivity of a biological surface through the sequential administration of the above-noted components whereby the resulting conjugate is bound to a natural or synthetic surface to enhance the acoustic reflectivity thereof for ultrasonic imaging. The invention is also directed to compositions for use in ultrasonic imaging of such surfaces and for enhancing the acoustic reflectivity thereof.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

**[0009]** Figure 1 is a graph showing changes in aggregate particle size of biotinylated and control perfluorocarbon emulsions with increasing avidin concentration;

**[0010]** Figure 2 shows ultrasonic images of control and biotinylated perfluorocarbon emulsion before and after the addition of avidin;

**[0011]** Figure 3 is a graphic illustration of dialysis tubing images and region of interest placement for gray scale analysis;

**[0012]** Figure 4 is a graph showing changes in average pixel gray scale associated with the addition of avidin to control or biotinylated perfluorocarbon emulsion;

**[0013]** Figure 5 is a graph showing the effect of control and biotinylated perfluorocarbon emulsion on apparent backscatter transfer function and integrated backscatter of avidinized nitrocellulose membranes;

**[0014]** Figure 6 is a graph showing the apparent backscatter transfer function of biotinylated and control perfluorocarbon emulsions targeted to D-dimer covalently conjugated to nitrocellulose membranes;

**[0015]** Figure 7 is a graph showing the apparent backscatter transfer function (dB) of biotinylated and control perfluorocarbon emulsions at low ultrasonic frequencies;

**[0016]** Figure 8 is a graph showing the apparent backscatter transfer function of biotinylated and control perfluorocarbon large particle size emulsions targeted to avidinized nitrocellulose membranes;

**[0017]** Figure 9 shows ultrasonic images of plasma thrombi before and after exposure to control or biotinylated emulsions;

**[0018]** Figure 10 is a graph showing the average pixel grayscale level of plasma thrombi pretargeted with antifibrin monoclonal antibody and exposed to control or biotinylated perfluorocarbon emulsion;

**[0019]** Figure 11 shows ultrasonic images of femoral artery thrombus acoustically enhanced with biotinylated perfluorocarbon emulsion in vivo;

**[0020]** Figure 12 is a graph showing the net change in apparent backscatter transfer function of biotinylated and control perfluorocarbon emulsions targeted to prostate specific antigen in prostatic carcinoma relative to normal regions;

**[0021]** Figure 13 is a graph showing the net change in integrated backscatter between normal prostatic stroma and cancer regions for control versus biotinylated perfluorocarbon emulsions;

**[0022]** Figure 14 is a graph showing the net change in apparent backscatter transfer function of biotinylated and control perfluorocarbon emulsions targeted to OC-125 antigen in ovarian carcinoma relative to normal regions;

**[0023]** Figure 15 is a graph showing the net change in integrated backscatter between normal ovarian tissue and carcinoma regions for control versus biotinylated perfluorocarbon emulsions;

**[0024]** Figure 16 shows ultrasonic and optical images of tonsil using perfluorocarbon contrast and horseradish peroxidase targeted to epithelium with anticytokeratin antibodies; and

**[0025]** Figure 17 shows peak detected ultrasonic images of tonsil epithelium acoustically enhanced with anticytokeratin antibody targeted perfluorocarbon emulsion.

**[0026]** Figure 18 is a graph showing backscattered power from plasma clots after the one step fibrin targeted emulsion, the backscatter being computed from the average of the frequency-dependent backscatter transfer function over the useful bandwidth of the transducer;

**[0027]** Figure 19 shows scanning electron microscope images of cell cultures treated with no (Fig. 19A) or untargeted control emulsion (Fig. 19B), tissue

factor targeted specifically bound to the aortic smooth muscle cells (Fig. 19C) and binding specifically blocked by anti-tissue factor antiserum (Fig. 19D);

**[0028]** Figure 20 shows ultrasonic images of vessels before and after treatment with targeted conjugated emulsion; and

**[0029]** Figure 21 shows the results of staining which revealed a pattern of localization of tissue factor in balloon-injured vessels in the media (arrows) at the site of acoustic contrast enhancement, as well as on the internal surface associated with inflammatory cells (Fig. 21 A; L-lumen) and control uninjured vessels exhibiting staining primarily on the adventitia and not in the tunica media (Fig. 21B).

### **DESCRIPTION OF THE PREFERRED EMBODIMENTS**

**[0030]** In accordance with the present invention, it has now been found that a ligand-based binding system having broad application may be achieved through ligand-based binding of lipid encapsulated particles to molecular epitopes on a surface in vivo or in vitro by sequentially administering (a) a site-specific ligand activated with a biotin activating agent; (b) an avidin activating agent; and (c) lipid encapsulated particles activated with a biotin activating agent, whereby the ligand is conjugated to the lipid encapsulated particles through an avidin-biotin interaction or complexing and the resulting conjugate is bound to the molecular epitopes on the surface. The ligand-based binding system of the present invention thus permits detection of molecular moieties such as peptides, carbohydrates or nucleic acids with a specific ligand probe (e.g. an antibody or antibody fragment) complexed or conjugated with avidin and biotin, the latter being carried by lipid encapsulated particles (e.g. biotinylated lipid encapsulated emulsion or liposome). The ligand-based binding system of the invention may be employed in an ultrasonic contrast agent system, ultrasound-based ELISA-like laboratory diagnostic assays in liquid and solid phase systems and in cell cultures, electrophoretic, chromatographic and hybridization detection systems, and for the detection of thrombi, infections, cancers and infarctions in patients with the use of conventional ultrasonic imaging methods. The invention may also be applied for therapeutic purposes by delivery of chemotherapeutic agents or drugs to desired sites due to the specificity of the binding system coupled with the ability to monitor the progress of the therapeutic treatment through repeated imaging at such sites. In this regard, the above-referred to conjugate of the ligand to the lipid encapsulated particles through an avidin-biotin interaction or complexing is effective

for imaging by x-ray, ultrasound, magnetic resonance or positron emission tomography.

**[0031]** In one embodiment of the invention, there is provided a method for enhancing the reflectivity of a biological surface by sequentially administering to the surface (a) a site-specific ligand activated with a biotin activating agent; (b) an avidin activating agent; and (c) lipid encapsulated particles activated with a biotin activating agent; whereby the ligand is conjugated to the lipid encapsulated particles through an avidin-biotin interaction and the resulting conjugate is bound to the biological surface to enhance the acoustic reflectivity thereof for ultrasonic imaging. This novel triphasic approach utilizes an avidin-biotin interaction to permit administration of the targeting ligand separate from the acoustic lipid encapsulated particles. In a specific application of the method in accordance with the invention, a biotinylated ligand is first systemically administered to a patient to pretarget the tissue or biological surface of interest and to circulate for a period of time necessary or sufficient to optimize the percentage bound. In the second phase, avidin is administered, circulates and binds to the biotinylated ligand attached to the target tissue or surface and to any residual, free circulating ligand. Avidin cross-linking increases the avidity and stability of the ligand on the target tissue or surface while promoting the rapid clearance of circulating avidin-ligand complexes via the reticuloendothelial system. In the third phase, the biotinylated lipid encapsulated particles are administered, binding to avidin through unoccupied biotin binding sites, and imparting increased acoustic contrast to the targeted tissue surface. Repeated sequential administration of avidin and the biotinylated lipid encapsulated particles may be carried out to amplify the acoustic contrast effect of the lipid encapsulated particles bound to the targeted surface.

**[0032]** In the practice of the invention, the ligand employed may be, for example, constituted by monoclonal or polyclonal antibodies, viruses, chemotherapeutic agents, receptor agonists and antagonists, antibody fragments, lectin, albumin, peptides, hormones, amino sugars, lipids, fatty acids, nucleic acids and cells prepared or isolated from natural or synthetic sources. In short, any site-specific ligand for any molecular epitope or receptor to be detected through the practice of the invention may be utilized.

**[0033]** The ligand is activated with a biotin activating agent. As employed herein, the term "biotin activating agent" or "biotinylated" encompasses

biotin, biocytin and other biotin analogs such as biotin amido caproate N-hydroxysuccinimide ester, biotin 4-amidobenzoic acid, biotinamide caproyl hydrazide and other biotin derivatives and conjugates. Other derivatives include biotin-dextran, biotin-disulfide-N-hydroxysuccinimide ester, biotin-6 amido quinoline, biotin hydrazide, d-biotin-N hydroxysuccinimide ester, biotin maleimide, d-biotin p-nitrophenyl ester, biotinylated nucleotides and biotinylated amino acids such as Nε-biotinyl-L-lysine.

**[0034]** In the second phase, as previously mentioned, an avidin activating agent is administered. As employed herein, the term "avidin activating agent" or "avidinized" encompasses avidin, streptavidin and other avidin analogs such as streptavidin or avidin conjugates, highly purified and fractionated species of avidin or streptavidin, and non or partial amino acid variants, recombinant or chemically synthesized avidin analogs with amino acid or chemical substitutions which still accommodate biotin binding.

**[0035]** The lipid encapsulated particles or contrast agent employed in the third phase may be constituted, for example, by a biotinylated emulsion or liposome which may contain a gas, liquid or solid. In a specific example, the lipid encapsulated particles may be constituted by a perfluorocarbon emulsion, the emulsion particles having incorporated into their outer coating a biotinylated lipid compatible moiety such as a derivatized natural or synthetic phospholipid, a fatty acid, cholesterol, lipolipid, sphingomyelin, tocopherol, glucolipid, stearylamine, cardiolipin, a lipid with ether or ester linked fatty acids or a polymerized lipid. Thus, the biotinylated contrast agent constituting the lipid encapsulated particles may be produced by incorporating biotinylated phosphatidylethanolamine into the outer lipid monolayer of a perfluorocarbon emulsion.

**[0036]** Perfluorocarbon emulsions are particularly well suited for biomedical applications and for use in the practice of the present invention. They are known to be stable, biologically inert and readily metabolized, primarily by trans-pulmonic alveolae evaporation. Further, their small particle size easily accommodate transpulmonic passage and their circulatory half-life (4-8 hours) advantageously exceeds that of other agents. Also, perfluorocarbons have been used to date in a wide variety of biomedical applications, including use as artificial blood substitutes. For use in the present invention, various fluorocarbon emulsions may be employed including those in which the fluorocarbon is a fluorocarbon-hydrocarbon, a



perfluoroalkylated ether, polyether or crown ether. Useful perfluorocarbon emulsions are disclosed in U.S. Patent Nos. 4,927,623, 5,077,036, 5,114,703, 5,171,755, 5,304,325, 5,350,571, 5,393,524, and 5,403,575 and include those in which the perfluorocarbon compound is perfluorotributylamine, perfluorodecalin, perfluorooctylbromide, perfluorodichlorooctane, perfluorodecane, perfluorotripropylamine, perfluorotrimethylcyclohexane or other perfluorocarbon compounds. Further, mixtures of such perfluorocarbon compounds may be incorporated in the emulsions utilized in the practice of the invention. As a specific example of a perfluorocarbon emulsion useful in the invention may be mentioned a perfluorodichlorooctane emulsion wherein the lipid coating thereof contains between approximately 50 to 99.5 mole percent lecithin, preferably approximately 55 to 70 to mole percent lecithin, 0 to 50 mole percent cholesterol, preferably approximately 25 to 45 mole percent cholesterol and approximately 0.5 to 10 mole percent biotinylated phosphatidylethanolamine, preferably approximately 1 to 5 mole percent biotinylated phosphatidylethanolamine. Other phospholipids such as phosphatidylserine may be biotinylated, fatty acyl groups such as stearylamine may be conjugated to biotin, or cholesterol or other fat soluble chemicals may be biotinylated and incorporated in the lipid coating for the lipid encapsulated particles. The preparation of an exemplary biotinylated perfluorocarbon for use in the practice of the invention is described hereinafter in accordance with known procedures.

**[0037]** When the lipid encapsulated particles are constituted by a liposome rather than an emulsion, such a liposome may be prepared as generally described in the literature (see, for example, Kimelberg et al., CRC Crit. Rev. Toxicol. 6,25 (1978) and Yatvin et al., Medical Physics, Vol. 9, No. 2, 149 (1982)). Liposomes are known to the art and generally comprise lipid materials including lecithin and sterols, egg phosphatidyl choline, egg phosphatidic acid, cholesterol and alpha-tocopherol.

**[0038]** With respect to the particle size of the lipid encapsulated particles constituted by a perfluorocarbon emulsion or liposome, the particle size may range between approximately 0.05 to 5 microns and preferably between approximately 0.05 and 0.5 micron. Small size particles are thus preferred because they circulate longer and tend to be more stable than larger particles.

**[0039]** As indicated, the ligand is conjugated to the lipid encapsulated particles or perfluorocarbon emulsion through an avidin-biotin interaction. The ligand

may also be conjugated to the emulsion directly or indirectly through intervening chemical groups or conjugated directly or indirectly to biotin or a biotin analog through intervening chemical groups such as an alkane spacer molecule or other hydrocarbon spacer. The use of spacer molecules between the ligand and biotin or between biotin and the emulsion is not required but aids in rendering the biotin more available for binding to avidin.

**[0040]** In accordance with the broadest aspect of the invention, it has been found that liquid perfluorocarbon emulsions have very poor intrinsic echogenicity when free in suspension, but when bound to a surface, they increase the acoustic reflectivity of the surface. By conjugating the ligand directly to the emulsion and binding the resulting ligand-emulsion conjugate to a surface, enhanced acoustic reflectivity is realized for ultrasonic imaging.

**[0041]** As previously mentioned, the emulsion or liposome constituting the lipid encapsulated particles or vesicles may contain a gas, liquid or solid. The gas may be nitrogen, oxygen, carbon dioxide or helium and may, for example, be evolved from the fluorocarbon component of the emulsions described above.

**[0042]** Alternatively, but less preferably, the ligand-based binding method of the invention may be carried out by sequentially administering a site-specific ligand activated with a biotin or avidin activating agent and lipid encapsulated particles activated with a biotin or avidin activating agent, a biotin activating agent being used where an avidin activating agent was employed in the first step and an avidin activating agent being used where a biotin activating agent was employed in the first step. The direct conjugation of the ligand to a perfluorocarbon emulsion, for example, is less preferable since it may accelerate in vivo clearance of the emulsion contrast agent.

**[0043]** In the practice of the invention, it has been unexpectedly found that the individual components of the ultrasonic contrast agents as described above are poorly reflective or have low echogenicity in the bloodstream but become highly reflective when the ligand-avidin-emulsion complex is formed in vivo at the desired site or biological surface and thereby substantially enhances the acoustic reflectivity thereof for ultrasonic imaging. This is in sharp contrast to previously known sonographic contrast agents which are inherently bright or of high reflectivity in the bloodstream. The improved acoustic reflectivity achieved through the present

invention provides the advantage of enhancing the signal-to-noise ratio because the background contrast from lipid encapsulated particles in the blood is minimal. Thus, the present invention offers an improved noninvasive method for forming an acoustic contrast agent which can be targeted in vitro or in vivo and which when bound to a specific desired site alters the acoustic reflectivity of a tissue surface or support media in a manner detectable with ultrasonic transducers suitable for biomedical and diagnostic applications within a frequency range of at least 5 to 50 MHz (nominal center frequencies may be wider ranging based on the knowledge that these are broad band transducers). The method of the invention advantageously provides a practical means for detecting any molecular epitope or receptor for which a biotinylated monoclonal antibody or other ligand is available without the need for use of ionizing radiation with or without associated invasive procedures in various clinical applications and while employing standard, commercially available ultrasonic technology.

**[0044]** The present invention does not employ ultrasonic contrast systems or agents to delineate blood flow as in the prior art but rather to detect physiologic and pathologic events by sensing the accumulation of the contrast agent at specific binding sites.

**[0045]** In the application of the invention to diagnostic assays such as ultrasound-based ELISA-type laboratory diagnostic assays in liquid and solid phase systems, the surface on which ligand-based binding of lipid encapsulated particles to molecular epitopes occurs may be, for example, nylon, nitrocellulose membranes or a gel as well as a biological surface.

**[0046]** The ligand-based binding system of the invention may also be applied to provide a chemotherapeutic agent or gene therapy delivery system combined with ultrasonic imaging. For example, chemotherapeutic agents or immune activating drugs such as tissue plasminogen activator, adriamycin, vincristine, urokinase, streptokinase, methotrexate, cytarabine, thioguanine, doxorubicin, 5-fluorouracil, cisplatin, etoposide, ifosfamide, asparaginase, deoxycoformycin, hexamethyl melamine and including radioactive agents may be incorporated in the lipid encapsulated particles and become part of the conjugate bound to a specific biological surface site for therapeutic action. The present invention would also advantageously permit the site to be ultrasonically imaged in order to monitor the progress of the therapy on the site and to make desired adjustments in the dosage of

therapeutic agent subsequently directed to the site. The invention thus provides a noninvasive means for the detection and therapeutic treatment of thrombi, infections, cancers and infarctions in patients while employing conventional ultrasonic imaging systems.

**[0047]** The following examples illustrate the practice of the invention.

#### EXAMPLE 1

**[0048]** The procedure for preparing a biotinylated lipid encapsulated perfluorodichlorooctane emulsion for use in ultrasound imaging is as follows.

**[0049]** The biotinylated lipid perfluorodichlorooctane (PFDCO) emulsion is comprised of the following components: PFDCO (40% v/v), safflower oil (2.0%w/v), a surfactant co-mixture (2.0% w/v) and glycerin (1.7% w/v). The surfactant co-mixture is composed of approximately 64 mole% lecithin, 35 mole % cholesterol and 1 mole % N-(6-biotinoyl)amino)hexanoyl)dipalmitoyl-L-alpha-phosphatidylethano-lamine. These components are weighed together into a test tube and dissolved in chloroform. The chloroform is stripped from the material and the resulting surfactant mixture is dried in a 50°C vacuum oven overnight. The co-mixture is dispersed into water by sonication resulting in a liposome suspension. The suspension is transferred into a 30 mL capacity blender cup (Dynamics Corporation of America, New Hartford, CT) along with the PFDCO and oil. The mixture is blended for 30-60 seconds to a pre-emulsion. The preemulsified sample is transferred to the reservoir of a microfluidizer, model S 110 (Microfluidics, Newton, MA), and emulsified for three minutes at 10,000 psi. To prevent the emulsion from heating excessively during homogenization, the shear valve and mixing coil of the microfluidizer are immersed in a room temperature water bath during processing. The final temperature of the emulsion is approximately 35°C. The finished emulsion is bottled in 10 mL serum vials, blanketed with nitrogen gas and sealed with stopper/crimp seal. The average particle size of the finished product, measured by a laser light scatter particle sizer (Brookhaven Instruments Corporation, Holtsville, N-Y), is 250 nm.

#### EXAMPLE 2

**[0050]** The incorporation of biotinylated phosphatidylethanolamine into the encapsulating lipid monolayer of perfluorocarbon emulsion is prepared as

described in Example one and demonstrated to increase aggregate particle size in the presence of titrated concentrations of avidin (Pierce, Rockford, IL 61105). An identically prepared control emulsion is prepared which incorporates nonbiotinylated phosphatidylethanolamine into the outer lipid monolayer of the perfluorocarbon emulsion. Avidin is resuspended in isotonic phosphate buffered saline (PBS, Fisher Inc., Fair Lawn, NJ). Within a polystyrene cuvette, a 3.0 ml reaction mixture is prepared containing PBS, biotinylated or control perfluorocarbon emulsion (20  $\mu$ l) and avidin, at 0.0, 0.5, 1.0, 1.5 or 2.0  $\mu$ g/ml. Contents are mixed by gentle inversion and react for thirty minutes at room temperature. Emulsion particle sizes are determined in triplicate with a Brookhaven BI-90 particle size analyzer (Holtsville, NY) at 37°C. Aggregate particle size of the biotinylated emulsion increased progressively from a baseline of  $263 \pm 2.54$  nm to greater than 2000 nm with increasing concentration of avidin (Figure 1). Marked flocculation and sedimentation are noted when avidin concentrations exceed 2.0  $\mu$ g/ml. The particle size of the control emulsion is  $234 \pm 3.81$  nm in diameter and addition of 2.0  $\mu$ g of avidin to the reaction mixture does not affect particle size. These results clearly demonstrate that the biotinylated phosphatidylethanolamine is incorporated and oriented appropriately into the outer lipid monolayer of the perfluorocarbon emulsion and that surface biotins are adequately available to avidin in the media. Multiple biotin binding sites on the avidin molecule as well as multiple biotin residues on the surface of the emulsion progresses towards a rapid complexing of particles in vitro.

### EXAMPLE 3

**[0051]** Biotinylated perfluorocarbon emulsion particles, approximately 250 nm in diameter, with low independent acoustic reflectivity, are complexed with avidin in solution which eventuates aggregation and enhances echogenicity. Biotinylated and control perfluorocarbon emulsion (200  $\mu$ l) prepared as described previously are diluted in PBS (15 ml) and placed within dialysis tubing (Spectra/Por 4, 25 mm, MWCO 12,000-14,000, Spectrum Medical Industries, Inc., Los Angeles, CA), ultrasonically imaged within a PBS water bath at room temperature using a 7.5 MHz focused transducer and a Hewlett Packard (HP) Sonos 2500 Phased Array Imaging System (Andover, MA). Real-time images are recorded to SVHS video tape for subsequent image analysis. Pixel grayscale and homogeneity are assessed on selected freeze-frame images using NIH Image 1.47 (National Institutes of Health).

Avidin (30  $\mu$ /ml) is added to each emulsion suspension, mixed by gentle inversion and allowed to complex for 30 minutes. The emulsion suspensions are optically opaque but ultrasonically undetected prior to the addition of avidin. Complexing of the biotinylated perfluorocarbon emulsion ensues rapidly with the addition of avidin and a white, flocculant precipitate soon appears. Avidin induces no changes in control emulsion suspension. Insonification of the suspensions reveals that the biotinylated perfluorocarbon emulsion particles opacify the dialysis tubing; whereas, the control particles are not appreciated acoustically (Figure 2). Gray scale echo intensity analysis of freeze-frame images of the control and biotinylated emulsion suspensions before and after avidin are summarized in Figures 3 and 4. The increased average grayscale level of the biotinylated emulsion ( $71.3 \pm 22.1$ ) suspension relative to its pre-avidin pixel gray scale level ( $2.2 \pm 4.4$ ) demonstrates the acoustic enhancement achieved. Average pixel gray scale levels of the control emulsion before ( $3 \pm 7.33$ ) and after ( $1.0 \pm 1.3$ ) avidin addition are similar. These results demonstrate the low acoustic reflectivity of the perfluorocarbon emulsion when imaged as independent particles in comparison with the enhanced echogenic nature of the aggregated biotinylated particles. The lack of acoustic change in the control emulsion suspension in the presence of avidin confirms the ligand specificity of the biotinylated emulsion.

#### EXAMPLE 4

**[0052]** Biotinylated perfluorocarbon emulsion, approximately 250 nm diameter, are specifically targeted to avidin, covalently bound to a modified nitrocellulose membrane and increases the acoustic reflectivity of the membrane surface at high ultrasonic frequencies (30 to 60 MHz). Briefly, nitrocellulose membranes (S+S NCTM, Schleicher & Schuell, Keane, NH) were conjugated to avidin using a diaminohexane (Sigma Chemical Co., St. Louis, MO) spacer and glutaraldehyde (Sigma Chemical Co., St. Louis, MO) activation as described by Måsson et al. (Electrophoresis 1993, 14, 860-865). Nitrocellulose discs (2 cm diameter) are soaked in 2.5% diaminohexane dissolved in deionized water for 60 minutes with constant slow rotary agitation. Membranes are washed with 1M acetic acid for 6-7 hours followed by an 18+ hour deionized water wash with constant agitation. The membranes are placed in 1% glutaraldehyde in 0.1 M sodium bicarbonate buffer, pH 10.0 for 15 minutes then washed for three hours with deionized water. Nitrocellulose membranes are stored and dried at 4°C until use;

storage does not exceed three days. Fifty (50)  $\mu\text{l}$  of avidin (250  $\mu\text{g}$ ) are spotted dropwise upon the center of six membranes with a microliter syringe and allowed to dry. Each membrane is extensively washed with a 0.1% Tween-20 (Sigma Chemical Co., St. Louis, MO) in PBS then placed in 3% bovine serum albumin (BSA, crystallized, Sigma Chemical Company, St. Louis, MO) dissolved in PBS-0.1% Tween-20 for 20 minutes to blockade nonspecific protein binding sites around the periphery of the disc. After the BSA blockade, each disc is extensively washed with PBS and placed in 300  $\mu\text{l}$  of either biotinylated or control perfluorocarbon emulsion suspended in 4 ml PBS for 20 minutes. Unbound emulsion is removed in serial PBS washes. Each disc is reexposed to avidin and control or biotinylated emulsion to ensure saturated coverage of the nitrocellulose surface. The nitrocellulose discs are washed and stored in PBS at 4°C until imaged with acoustic microscopy.

**[0053]** For acoustic microscopic imaging, each nitrocellulose disc is placed flat above a polished stainless steel plate in a polystyrene holder with a 2 X 2 cm central window removed. The mounted specimen is immersed into PBS at ambient temperature for ultrasonic insonification. A custom designed acoustic microscope, utilizing a 50MHz (nominal frequency) broadband, focused, piezoelectric delay-line transducer (1/4 inch diameter, 1/2 inch focal length, Model V390, Panametrics Co., Waltham, MA) operated in the pulse-echo mode is utilized for insonification. Backscattered radio frequency (RF) data is collected and digitized at 500 megasamples per second utilizing a Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data are acquired from approximately 100 independent sites from each region of interest with 100 micron lateral step resolution.

**[0054]** A radio frequency peak-detected scan of the data is converted into a gray scale (0=lowest scattering, 255=highest scattering) map to allow selection of regions of interest for integrated backscatter analysis. Radio frequency (RF) ultrasonic data are stored in a raster scan format and analyzed with custom software. Segments of the RF lines are gated for integrated backscatter analysis to encompass the front and back surfaces of the nitrocellulose disc. The data are multiplied by a rectangular window and their power spectra are determined by fast-Fourier transformation. The power spectra from the specimens referenced to the power spectrum returned from a near-perfect steel planar reflector and the frequency

dependent backscatter transfer function across the useful bandwidth of the transducer (30 to 60 MHz) are computed and expressed in decibels relative to acoustic scattering from the near perfect steel plate reflector (Wong et al., *Ultrasound in Med & Biol.* 1993; 19: 365-374). Integrated backscatter (IB) is computed as the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer.

**[0055]** Discs incubated with biotinylated perfluorocarbon emulsion have central regions with high acoustic scattering in comparison with the peripheral (i.e. background) regions of the same disc. Nitrocellulose discs incubated with the control emulsion have no central high scattering regions and no differences in acoustic character is detected by changes in the RF signature between the central and peripheral regions of the disc. IB from the centrally located, biotinylated emulsion region ( $-17.8 \pm 0.2$  db) is  $6.3 \pm 0.1$  dB(4-fold) greater ( $p < 0.05$ ) than IB from the analogous region on the control disc ( $-24.1 \pm 0.2$  dB). The frequency-dependent variation in apparent backscatter transfer function (mean  $\pm$  SEM) from the avidin spotted regions of the biotinylated and control emulsion discs are presented in Figure 5. A smooth and consistently greater acoustic response is noted across the frequency spectrum due to the bound biotinylated emulsion. These results demonstrate the effectiveness of the biotinylated perfluorocarbon emulsion to specifically target a surface bound antigen and dramatically alter the acoustic reflectivity of the surface with the bathing medium, increasing the ultrasonic backscattered power at high frequencies.

#### EXAMPLE 5

**[0056]** Biotinylated perfluorocarbon emulsion (250 nm diameter) is specifically targeted to D-dimer covalently attached to a modified nitrocellulose membrane utilizing a biotinylated anti-D-dimer  $F_{(ab)}$  fragment-avidin complex and results in a marked increase in the acoustic power reflected from the surface. D-dimer is covalently linked to nitrocellulose discs modified with a diaminohexane spacer arm and activated with glutaraldehyde as previously described in Example 4. Fifty (50)  $\mu$ g of D-dimer is spotted with a microliter syringe upon the center of three of six membranes and allowed to air dry. Unbound D-dimer is exhaustively washed from the membranes with phosphate buffered saline (PBS)-0.1 % Tween-20. Nonspecific protein binding sites of all membranes are blocked with 3% bovine serum albumin



(BSA) in PBS-0. 1% Tween-20 for 20 minutes followed by serial PBS washes. D-dimer spotted membranes are incubated with 12.5  $\mu$ g biotinylated anti-D-dimer F(ab) antibody in 4.0 ml 3% BSA for 2 hours, washed with PBS buffer and then incubated with 250  $\mu$ g avidin in 4 ml PBS for 30 min. After removing unbound avidin with PBS washes, the discs are exposed to either biotinylated or control perfluorocarbon emulsion (300  $\mu$ l) in 4.0 ml PBS for 20 minutes. Excess emulsion is removed with PBS buffer washes. Discs are reexposed to avidin and perfluorocarbon emulsion as described above and the membranes are stored in PBS at 4°C until imaging.

**[0057]** For acoustic microscopic imaging, each nitrocellulose disc is placed flat above a polished stainless steel plate in a polystyrene holder, immersed in PBS at ambient temperature, and insonified with a custom designed acoustic microscope, utilizing a 50MHz (nominal frequency) broadband, focused, piezoelectric delay-line transducer (1/4 inch diameter, 1/2 inch focal length, Model V390, Panametrics Co., Waltham, MA) operated in the pulse-echo mode. Backscattered radio frequency (R.F) data is collected and digitized at 500 megasamples per second utilizing a Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data are acquired from approximately 100 independent sites from each region of interest with 100 micron lateral step resolution.

**[0058]** A radio frequency peak-detected scan of the data is converted into a gray scale (0=lowest scattering, 255=highest scattering) map of the disc to allow visual inspection and selection of regions of interest for integrated backscatter (IB) analysis. Radio frequency (RF) ultrasonic data are stored in a raster scan format and analyzed with custom software. Segments of the RF lines are gated for integrated backscatter analysis to encompass the front and back surfaces of the nitrocellulose disc. The data are multiplied by a rectangular window and their power spectra are determined by fast-Fourier transformation. The power spectra from the specimens referenced to the power spectrum returned from a near-perfect steel planar reflector and the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer (30 to 60 MHz) are computed and expressed in decibels relative to acoustic scattering from the near perfect steel plate reflector (Wong et al., *Ultrasound in Med & Biol.* 1993; 19: 365-374). Integrated backscatter is computed as

the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer.

**[0059]** Biotinylated, anti-D-dimer F(ab) fragment is specifically bound to the central region of the D-dimer spotted discs and crosslinked by avidin through its biotin moiety. As in previous examples, biotinylated perfluorocarbon emulsion specifically binds to the antibody bound avidin; whereas, the nonspecific binding of the control emulsion have no binding and are not detected acoustically. IB of the biotinylated emulsion coated nitrocellulose ( $-18.0 \pm 0.2$  dB) was greater by  $4.6 \pm 0.1$  dB ( $p < 0.05$ ) than that from the control disc ( $-22.6 \pm 0.1$  dB) over the 30 to 60 MHz frequency range. The frequency-dependent variation in apparent backscatter transfer function (mean  $\pm$  SEM) of the biotinylated and control emulsion discs are presented in Figure 6. A smooth and consistently greater acoustic response is noted across the frequency spectrum due to the bound biotinylated emulsion. These data confirm and extend the findings of Example 4 with avidin alone, demonstrating that biotinylated perfluorocarbon emulsion bound through a specific, targeting ligand system can significantly enhance the acoustic backscatter of a solid support surface.

#### EXAMPLE 6

**[0060]** Biotinylated perfluorocarbon emulsion (250 nm diameter) is specifically targeted to avidin conjugated to nitrocellulose discs and insonified at clinically relevant frequencies (5 to 15 MHz) and significantly increases the acoustic backscatter of the membrane. Briefly, nitrocellulose membranes (S+S NCTM, Schleicher & Schuell, Keane, NH) are conjugated to avidin using a diaminohexane (Sigma Chemical Co., St. Louis, MO) spacer and glutaraldehyde (Sigma Chemical Co., St. Louis, MO) activation as described by Másson et al. (Electrophoresis 1993, 14, 860-865). Nitrocellulose discs (2 cm diameter) are soaked in 2.5% diaminohexane dissolved in deionized water for 60 minutes with constant, slow rotary agitation. Membranes are transferred to and washed with 1M acetic acid for 6-7 hours then transferred for continued washing in deionized water for at least 18 additional hours with constant agitation. The membranes are placed in 1% glutaraldehyde in 0.1M sodium bicarbonate buffer, pH 10.0 for 15 minutes. After glutaraldehyde activation is complete, the membranes are washed with continued agitation for three hours. The nitrocellulose membranes stored and dried at 4°C until use; storage does not exceed three days. Fifty (50)  $\mu$ l of avidin (250  $\mu$ g) are spotted dropwise upon the center of a

nitrocellulose membrane with a microliter syringe and allowed to dry. Each membrane is washed with 0.1 % Tween-20 (Sigma Chemical Co., St. Louis, MO) in phosphate buffered saline (PBS) then placed in 3% bovine serum albumin (BSA, crystallized, Sigma Chemical Company, St. Louis, MO) dissolved PBS-0.1 % Tween-20 for 20 minutes to blockade nonspecific protein binding sites around the periphery of the disc. After the BSA blockade, each disc is washed with PBS and placed in 300  $\mu$ l of either biotinylated or control perfluorocarbon emulsions suspended in 4 ml PBS for 20 minutes with mild, rotary agitation. The unbound emulsion is removed with washes of PBS. Each disc is reexposed to avidin, washed with PBS, reexposed to control or biotinylated perfluorocarbon emulsion and rewashed with PBS as previously described. The nitrocellulose discs are stored in PBS at 4°C until imaged with the acoustic microscope.

**[0061]** For acoustic microscopic imaging, each nitrocellulose disc is placed flat above a polished stainless steel plate in a polystyrene holder with a 2 cm x 2 cm central window removed. The mounted specimen is immersed into PBS at ambient temperature for ultrasonic insonification. A custom designed acoustic microscope, utilizing a 10 MHz (nominal frequency) broadband, focused, piezoelectric delay-line transducer (1/2 inch diameter, 2 inch focal length, Model V31 1, Panametrics Co., Waltham, MA) operated in the pulse-echo mode is utilized for insonification. Backscattered radio frequency (RF) data is collected and digitized at 500 megasamples per second utilizing a Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data are acquired from approximately 100 independent sites from each region of interest with 250 micron lateral step resolution. A radio frequency peak-detected scan of the data is converted into a gray scale (0=lowest scattering, 255=highest scattering) map of the disc to allow visual inspection and selection of regions of interest for integrated backscatter analysis. Radio frequency ultrasonic data are stored in a raster scan format and analyzed with custom software. Segments of the RF lines are gated for integrated backscatter analysis to encompass the front and back surfaces of the nitrocellulose disc. The data are multiplied by a rectangular window and their power spectra are determined by fast-Fourier transformation. The power spectra from the specimens referenced to the power spectrum returned from a near-perfect steel planar reflector and the frequency-dependent backscatter transfer function across the useful

bandwidth of the transducer (5 to 15 MHz) are computed and expressed in decibels relative to acoustic scattering from the near perfect steel plate reflector (Wong et al., *Ultrasound in Med & Biol.* 1993; 19: 365-374). Integrated backscatter (IB) is computed as the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer.

**[0062]** Discs incubated with biotinylated perfluorocarbon emulsion have central regions with high acoustic scattering relative to the peripheral regions of the same disc or the central regions of the control emulsion disc. Nitrocellulose discs incubated with the control emulsion have no high scattering regions. IB of the biotinylated emulsion coated nitrocellulose ( $0.5 \pm 0.5$  dB) was greater by  $9.6 \pm 0.1$  dB (8-fold) ( $p < 0.05$ ) than that from the control disc ( $-9.2 \pm 0.5$  dB) over the 5 to 15 MHz frequency range. The frequency-dependent variation in apparent backscatter transfer function (mean  $\pm$  SEM) of the biotinylated and control emulsion discs are presented in Figure 7. A smooth and consistently greater acoustic response is noted across the frequency spectrum due to the bound biotinylated emulsion. These data confirm and extend the findings of Examples 4 and 5 with avidin and D-dimer, demonstrating that biotinylated perfluorocarbon emulsion bound through a specific, targeting ligand system can significantly enhance the acoustic backscatter of a solid support surface and that this improved acoustic backscatter is detected at low, clinically useful ultrasonic frequencies (5 to 15 MHz) as well as high frequencies (30 to 60 MHz).

#### EXAMPLE 7

**[0063]** Biotinylated perfluorocarbon contrast, approximately 3000 nm diameter, is specifically targeted to avidin conjugated to nitrocellulose discs and insonified at clinically relevant frequencies (at least 5 to 15 MHz bandwidth). Briefly, nitrocellulose membranes (S+S NCTM, Schleicher & Schuell, Keane, NH) are conjugated to avidin using a diaminohexane (Sigma Chemical Co., St. Louis, MO) spacer and glutaraldehyde (Sigma Chemical Co., St. Louis, MO) activation as described by Másson et al. (*Electrophoresis* 1993, 14, 860-865). Nitrocellulose discs (2 cm diameter) are soaked in 2.5% diaminohexane dissolved in deionized water for 60 minutes with constant, slow rotary agitation. Membranes are transferred to and washed with 1M acetic acid for 6-7 hours then transferred for continued washing in deionized water for at least 18 additional hours with constant agitation. The membranes are placed in 1% glutaraldehyde in 0.1 M sodium bicarbonate buffer, pH

10. 0 for 15 minutes. After glutaraldehyde activation is complete, the membranes are washed with continued agitation for three hours. The nitrocellulose membranes stored and dried at 4°C until use; storage does not exceed three days.

**[0064]** Fifty (50)  $\mu$ l of avidin (250  $\mu$ g) are spotted dropwise upon the center of two of four membranes with a microliter syringe and allowed to dry. Each membrane is washed with 0.1 % Tween-20 (Sigma Chemical Co., St. Louis, MO) in phosphate buffered saline (PBS) then placed in 3% bovine serum albumin (BSA, crystallized, Sigma Chemical Company, St. Louis, MO) dissolved PBS-0.1 % Tween-20 for 20 minutes to blockade nonspecific protein binding sites around the periphery of the disc. After the BSA blockade, each disc is washed with PBS and placed in 300  $\mu$ l of either biotinylated or control perfluorocarbon emulsions, approximately 3000 nm particle size, suspended in 4 ml PBS for 20 minutes with mild, rotary agitation. The unbound emulsion is removed with washes of PBS. Each disc is reexposed to avidin, washed with PBS, exposed to perfluorocarbon emulsion and rewashed with PBS as previously described. The nitrocellulose discs are stored in PBS at 4°C until imaged with the acoustic microscope.

**[0065]** For acoustic microscope imaging, each nitrocellulose disc is placed flat above a polished stainless steel plate in a polystyrene holder with a 2 cm x 2 cm central window removed. The mounted specimen is immersed into PBS at ambient temperature for ultrasonic insonification. A custom designed acoustic microscope, utilizing a broadband 10 MHz (nominal frequency) focused, piezoelectric delay-line transducer (1/2 inch diameter, 2 inch focal length, Model V31 1, Panametrics Co., Waltham, AM) operated in the pulse-echo mode is utilized for insonification. Backscattered radio frequency (RF) data is collected and digitized at 500 megasamples per second utilizing a Tektronic DSA 601 digitizing oscilloscope (Beaverton, OR) with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data are acquired from approximately 100 independent sites from each region of interest with 250 micron lateral step resolution.

**[0066]** A radio frequency peak-detected scan of the data is converted into a gray scale (0=lowest scattering, 255=highest scattering) map of the disc to allow visual inspection and selection of regions of interest for integrated backscatter analysis. Radio frequency ultrasonic data are stored in a raster scan format and analyzed with custom software. Segments of the RF lines are gated for integrated

backscatter (IB) analysis to encompass the front and back surfaces of the nitrocellulose disc. The data are multiplied by a rectangular window and their power spectra are determined by fast-Fourier transformation. The power spectra from the specimens referenced to the power spectrum returned from a near-perfect steel planar reflector and the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer (5 to 15 MHz) are computed and expressed in decibels relative to acoustic scattering from the near perfect steel plate reflector (Wong et al., *Ultrasound in Med. & Biol.* 1993; 19: 365-374). Integrated backscatter is computed as the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer.

**[0067]** Discs incubated with biotinylated perfluorocarbon emulsion have central regions with high acoustic scattering relative to the peripheral regions of the same disc and central regions of the control disc. Nitrocellulose discs incubated with the control emulsion have no central high scattering regions and no differences in acoustic character are detected between the central and peripheral regions of the disc. IB of the biotinylated emulsion coated nitrocellulose ( $-2.4 \pm 0.7$  dB) was greater by  $8.8 \pm 0.3$  dB (approximately 8-fold ( $p < 0.05$ )) than that from the control disc ( $-11.2 \pm 0.4$  dB) over the 5 to 15 MHz frequency range. The frequency-dependent variation in apparent backscatter transfer function (mean  $\pm$  SEM) of the biotinylated and control emulsion discs are presented in Figure 8. A smooth and consistently greater acoustic response is noted across the frequency spectrum due to the bound biotinylated emulsion. These data confirm and extend the findings of Examples 4, 5 and 6 with avidin and D-dimer, demonstrating that biotinylated perfluorocarbon emulsions with large particle sizes can be bound through a specific, targeting ligand system and significantly enhance the acoustic backscatter of a solid support surface. This improved acoustic backscatter is detected at clinically relevant ultrasonic frequencies, 5 to 15 MHz.

#### EXAMPLE 8

**[0068]** Biotinylated perfluorocarbon emulsion is targeted to a plasma thrombi using biotinylated antifibrin monoclonal antibodies (NIBIHIO; Tymkewycz et al. 1993. *Blood Coagulation and Fibrinolysis* 4:211-221) and avidin. In a representative study (1 of 5), whole porcine blood is obtained and anticoagulated (9:1, v/v) with sterile sodium citrate. Blood is centrifuged at 1500 RPM at room

temperature and the plasma fraction is obtained and stored at 4°C. Two porcine plasma thrombi are produced by combining plasma, 100 MM calcium chloride (3:1 v/v) and 2-5 U thrombin in a plastic tube through which 5-0 Vicryl suture is passed. Thrombi are allowed to coagulate at room temperature.

**[0069]** One thrombus is incubated with 150 ug antifibrin monoclonal antibody in 10 ml PBS with 1% bovine serum albumin (BSA) for two hours and a second control thrombus is incubated in PBS with 1% BSA. The antibody treated thrombus is then incubated with 0.5 mg avidin in 10 ml PBS with 1% BSA for 30 minutes. The control thrombus remains in PBS with 1% BSA. Both thrombi are washed extensively with PBS. Each thrombus is incubated with 300 µl/10 ml PBS of either biotinylated or control emulsion for 30 minutes. All thrombi are reexposed to emulsion twice to ensure uniform coverage and ultrasonically insonified (Figure 9). Ultrasonic imaging is performed using a 7.5 MHz focused, linear phased array transducer and a Hewlett Packard Sonos 2500 Imaging System (Hewlett Packard, Inc., Andover, MA). All ultrasonic recordings are produced with fixed gain, compensation and time-gain compensation levels and are recorded on to SVHS videotape for subsequent image analysis. Average pixel grayscale over an extensive region of interest was sampled for 21 independent freeze-frame images for each thrombus using NIH Image 1.47 (National Institutes of Health; Figure 10). The biotinylated perfluorocarbon emulsion is found to provide a marked acoustic enhancement of the surface. Average pixel grayscale levels of the biotinylated emulsion thrombus are  $79.5 \pm 2.5$  whereas the brightness of the control was markedly less ( $34.8 \pm 2.2$ ,  $p < 0.05$ ). These results demonstrate the ability of biotinylated perfluorocarbon emulsion to target and acoustically enhance a biological tissue (i.e. thrombus) in vitro.

#### EXAMPLE 9

**[0070]** Biotinylated perfluorocarbon emulsion is targeted via biotinylated antifibrin antibodies (NIB5F3 and NIBIH10 Tynikewycz et al. 1993. Blood Coagulation and Fibrinolysis 4:211-221) to an isolated femoral artery thrombus in six mongrel dogs. A mongrel dog is anesthetized with sodium pentobarbital induction and halothane anesthesia. The right femoral artery and all branches are isolated at the level of the saphenous branch. A silver plated copper wire attached to a 22 ga. right angled needle point, insulated with plastic tubing (polyethylene P-240), is

inserted into the femoral artery and secured with 4-0 Prolene suture. A current of 200-400  $\mu$ A is applied for up to two hours. Thrombus formation is monitored with continuous wave doppler and discontinued after an approximately 50% increase in circulation velocity is noted distal to the electrical injury. Adventitial discoloration secondary to the current is appreciated proximal to the entry point of the wire. A 20 ga. catheter is inserted into a proximal branch of the femoral artery and secured with 4-0 silk suture. A pressurized 0.9% NaCl drip is attached through a three-way stopcock to the catheter. Blood flow into the isolated segment is disrupted by proximal snare ligature. Excess blood is flushed from the arterial segment to inhibit further thrombus formation by infusion of saline for 15 minutes. The distal draining branches of the femoral artery are ligated or snared with suture. Biotinylated antifibrin monoclonal antibody (50  $\mu$ g/1.0 ml PBS) is injected via the catheter and flushed with a few drops of saline. The antibody is allowed to incubate for one hour then the snare ligature distal to the wire insertion is released and excess antibody is flushed through with saline for five minutes. The distal femoral artery is reoccluded and avidin (250  $\mu$ g/1.0 ml PBS) is infused and incubates for 30 minutes. The distal ligature is again released and excess avidin is flushed through with saline for five minutes. The distal ligature is reestablished and biotinylated perfluorocarbon emulsion is infused and incubates for 30 minutes. After the initial exposure of the thrombus to the emulsion, the unbound emulsion is washed through with saline. Thrombi are each exposed to avidin and biotinylated perfluorocarbon emulsion as described above. In three animals, the contra lateral artery is also isolated, partially occluded with electrically induced thrombi and exposed to a control perfluorocarbon emulsion analagous to the administration of biotinylated emulsion described above. Femoral arteries exposed to either control or biotinylated perfluorocarbon emulsion are ultrasonically imaged at 7.5 MHz with a focused, linear phased array transducer and a clinical Hewlett-Packard Sonos 2500 Ultrasonic Imaging System before and after contrast administration. Acutely formed thrombi, both control and contrast targeted, are not ultrasonically appreciated. For 6 of 6 femoral arteries, partially occlusive thrombi are markedly enhanced using the antifibrin targeted biotinylated perfluorocarbon contrast. In 3 of 3 femoral arteries thrombi, exposure to the control perfluorocarbon emulsion does not accentuate their acoustic reflectivity and these thrombi remain ultrasonically undetectable. Figure 11 reveals a representative example of a femoral artery site of thrombus formation after electrical induction before and after exposure to antifibrin



antibody and biotinylated contrast. In the pre-contrast image, the femoral artery is observed with a bright echogenic wire point anode protruding into the lumen but no thrombus is appreciated. After treatment with the biotinylated contrast emulsion, a large partially occluded thrombus is clearly noted by the enhanced acoustic reflectivity (Figure 11). Again, no thrombus is appreciated in the control artery before or after exposure to control emulsion. These results demonstrate the concept of using bound perfluorocarbon emulsion to acoustically enhance biological surfaces, such as thrombotic tissue, in vivo to enable detection with a commercially available ultrasound imaging system.

#### EXAMPLE 10

**[0071]** Biotinylated perfluorocarbon emulsion, approximately 250 nm diameter, is targeted to prostatic carcinoma using monoclonal antibodies specific for prostate specific antigen (PSA) and are acoustically detected using polar, high frequency, high resolution acoustic microscopy. Representative examples of human prostatic carcinoma tissues are routinely processed by immersion fixation in 10% neutral buffered formalin and embedded in paraffin. Twenty micron sections are prepared for acoustic microscopy; 5 micron sections are used for optical studies. All histologic sections are mounted on acid cleaned glass slides that have been coated with poly-L-lysine. All mounted sections are heated at 55°C for 1 hour in an oven.

**[0072]** Prior to immunostaining, all sections are dewaxed in three changes of Americlear, and dehydrated in successive changes of 95% and 100% ethanol. Endogenous peroxidase activity is blocked only in sections prepared for optical studies by immersion in absolute methanol containing 0.6% (v/v) hydrogen peroxide for 30 minutes. These and all sections for acoustic microscopy are then rehydrated through graded ethanols and distilled water and placed in isotonic PBS (pH 7.4). All sections are incubated with target specific monoclonal antibodies.

**[0073]** Prostate sections are incubated with anti-PSA primary monoclonal antibodies per the recommendations of the vendor for 18 hours at 4°C in moisture chambers. After primary incubation, sections are rinsed in isotonic PBS, then overlain with a polyclonal biotinyl-horse anti-mouse immunoglobulin (VectaStain Elite Kits, Vector Laboratories, Burlingame, CA) for 1 hour at room temperature. After rinsing in PBS, a 30 micron section is prepared for acoustic microscopy. A section for light microscopy (5 micron) is incubated with avidin biotin-peroxidase

complex (VectaStain Elite Kit, Vector Lab) for 1 hour at room temperature. This section is rinsed in phosphate buffer (pH 7.6) and immersed in a solution of 3,3'-diaminobenzidine tetrahydrochloride (Sigma Chemicals, St. Louis, MO; 0.5 mg/ml in phosphate buffer, pH 7.6, containing 0.003% [v/v] hydrogen peroxide) for approximately ten minutes. The chromogenic precipitate is optically enhanced by brief immersion of stained sections in 0.125% (w/v) osmium tetroxide. The section is then rinsed in tap water, counterstained in Harris' hematoxylin, dehydrated in graded ethanols and Americlear, and mounted in a synthetic mounting medium.

**[0074]** After the second biotinylated antibody is incubated and washed, slides for acoustic microscopy are incubated in avidin (1.0 mg/-20cc PBS) using a bath on a rotating table for 30 min. Excess avidin is washed away with isotonic PBS buffer, pH 7.4-7.5 in three minute washes. Slides are incubated with biotinylated or control perfluorocarbon emulsion for twenty minutes (0.5 cc/-20.0 ml PBS), washed briefly with isotonic PBS 3X for 5 minutes each and reincubated with avidin (1.0 mg/-20cc) for 15 minutes. Excess avidin is rinsed off with three, 5 min. washes in PBS. The slide is then reincubated at above concentrations with biotinylated or control perfluorocarbon emulsion for 20 minutes. Unbound emulsion is washed away in three changes of PBS (5 minutes each) and the slides are transferred to the acoustic microscope for analysis.

**[0075]** The mounted specimens are each immersed into isotonic, phosphate buffered saline at room temperature for ultrasonic insonification. A custom designed acoustic microscope is used to collect ultrasonic data. The microscope consists of 50 MHz broadband, focused, piezoelectric delay-line transducer (1/4 inch diameter, 1/2 inch focal length, 62 micron beam diameter, Model V390, Panametrics Co., Waltham, MA) operated in the pulse-echo mode. A Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) is used to digitize 35 degree polar backscattered radio frequency (rf) data at 500 megasamples per second with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data is acquired from approximately 100 independent sites from each specimen with 50 micron lateral step resolution.

**[0076]** The rf data is stored in a low resolution raster scan format and analyzed with custom software. Segments of the rf lines are gated for integrated backscatter analysis to encompass the front surface (i.e. excluding the back wall). The gated data are multiplied by a Hamming window and their power spectra are

determined by fast-Fourier transformation. Power spectra within a tissue section are compared directly without reference to a steel plate. Integrated backscatter (IB) is computed from the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer (30 to 55 MHz). Immunostained tissues are reviewed using a Nikon Optiphot-2 microscope for regions of PSA positive staining and the acoustic characteristics are compared. The net change in the apparent backscatter transfer function between the normal prostatic stroma and carcinomatous regions are clearly increased in sections treated with PSA targeted biotinylated versus the control perfluorocarbon emulsion across the frequency spectrum (30 to 55 MHz; Figure 12). Biotinylated perfluorocarbon emulsion increases ( $p<0.05$ ) the integrated backscatter from regions of prostatic cancer ( $47.17\pm\text{dB}$ ) versus normal stromal ( $40.79\pm 1.18\text{ dB}$ ) by 6.38 dB (approximately 4-fold). In the control tissue sections, the integrated backscatter from the region of prostatic carcinoma ( $39.63\pm 1.63\text{ dB}$ ) was greater ( $p<0.05$ ) than that from the normal stromal areas ( $36.13\pm 2.17\text{ dB}$ ) by approximately 3.5 dB (2-fold), reflecting inherent differences in acoustic character between normal and cancerous prostatic tissue. However, the targeted biotinylated perfluorocarbon emulsion amplified ( $p<0.05$ ) these inherent differences by approximately 2-fold (2.87 dB; Figure 13). These results clearly demonstrate the ability of site-targeted biotinylated perfluorocarbon emulsion to specifically enhance acoustic detection of prostate cancer in vitro.

#### EXAMPLE 11

**[0077]** Biotinylated perfluorocarbon emulsion, approximately 250 nm diameter, is targeted to ovarian carcinoma using monoclonal antibodies specific for OC-125 antigen and are acoustically detected using polar, high frequency, high resolution acoustic microscopy. Representative examples of human ovarian, carcinoma tissues are routinely processed by immersion fixation in 10% neutral buffered formalin and embedded in paraffin. Twenty micron sections are prepared for acoustic microscopy; 5 micron sections are used for optical studies. All histologic sections are mounted on acid cleaned glass slides that have been coated with poly-L-lysine. All mounted sections are heated to 55°C for 1 hour in an oven.

**[0078]** Prior to immunostaining, all sections are dewaxed in three changes of Americlear, and dehydrated in successive changes of 95% and 100% ethanol. Endogenous peroxidase activity is blocked only in sections prepared for

optical studies by immersion in absolute methanol containing 0.6% (v/v) hydrogen peroxide for 30 minutes. These and all sections for acoustic microscopy are then rehydrated through graded ethanols and distilled water and placed in isotonic PBS (pH 7.4). All sections are incubated with target specific monoclonal antibodies.

**[0079]** Ovarian sections are incubated with anti-OC-125 primary monoclonal antibodies per the recommendations of the vendor for 18 hours at 4°C in moisture chambers. After primary incubation, sections are rinsed in isotonic PBS, then overlain with a polyclonal biotinyl-horse anti-mouse immunoglobulin (VectaStain Elite Kits, Vector Laboratories, Burlingame, CA) for 1 hour at room temperature. After rinsing in PBS, duplicate 30 micron sections are prepared for acoustic microscopy. Sections for light microscopy (5 micron) are incubated with avidin biotin-peroxidase complex (VectaStain Elite Kit, Vector Lab) for 1 hour at room temperature. Sections are rinsed in phosphate buffer (pH 7.6) and immersed in a solution at 3,3' diaminobenzidine tetrahydrochloride (Sigma Chemicals, St. Louis, MO; 0.5 mg/ml in phosphate buffer, pH 7.6, containing 0.0003% [v/v] hydrogen peroxide) for approximately ten minutes. The chromogenic precipitate is optically enhanced by brief immersion of stained sections in 0.125% (w/v) osmium tetroxide. Sections are then rinsed in tap water, counterstained in Harris' hematoxylin, dehydrated in graded ethanols and Americlear, and mounted in a synthetic mounting medium.

**[0080]** After the second biotinylated antibody is incubated and washed, slides are incubated in avidin (1.0 mg/-20 cc PBS) using a bath on a rotating table for 30 min. Excess avidin is washed away with isotonic PBS buffer, pH 7.4-7.5 in three 5 minute washes. The prepared slides are incubated with biotinylated or control perfluorocarbon emulsion for twenty minutes (0.5 cc/-20.0 ml PBS), washed briefly with isotonic PBS 3X for 5 minutes each and rewashed in avidin (1.0 mg/-20 cc) for 15 minutes. Excess avidin is rinsed off with three, 5 min. washes in PBS. Slides are then reincubated at above concentrations with biotinylated or control perfluorocarbon emulsion for 20 minutes. Unbound emulsion is washed away in three changes of PBS (5 minutes each) and the slides are transferred to the acoustic microscope for analysis.

**[0081]** The mounted specimens are each immersed into isotonic, phosphate buffered saline at room temperature for ultrasonic insonification. A custom designed acoustic microscope is used to collect ultrasonic data. The microscope

consists of 50MHz broadband, focused, piezoelectric delay-line transducer (1/4 inch diameter, 1/2 inch focal length, 62 micron beam diameter, Model V390, Panametrics Co., Waltham, MA) operated in the pulse-echo mode. A Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) is used to digitize 35 degree polar backscattered radio frequency (rf) data at 500 megasamples per second with 8-bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data is acquired from approximately 100 independent sites from each specimen with 50 micron lateral step resolution.

**[0082]** The rf data are stored in a low resolution raster scan format and analyzed with custom software. Segments of the rf lines are gated for integrated backscatter analysis to encompass the front surface (i.e. excluding the back wall). The gated data are multiplied by a Hamming window and their power spectra are determined by fast-Fourier transformation. Integrated backscatter is computed from the average of the frequency-dependent backscatter transfer function across the useful bandwidth of the transducer (30 to 55 MHz). The power spectra from the specimens are referenced to the power spectrum returned from a glass microscope slide. IB is expressed in decibels relative to the scattering from the glass slide. Immunostained tissues are reviewed using a Nikon Optiphot-2 microscope for regions of PSA positive staining and the acoustic characteristics are compared.

**[0083]** The net change in the apparent backscatter transfer function between the normal ovarian stroma and carcinomatous regions are clearly increased in sections treated with OC-125 targeted biotinylated versus the control perfluorocarbon emulsion across the frequency spectrum (30 to 55 MHz; Figure 14). Biotinylated perfluorocarbon emulsion increases ( $p < 0.05$ ) the integrated backscatter from regions of ovarian cancer ( $-28.19 \pm 1.39$  dB) versus normal stromal ( $-38.75 \pm 0.84$  dB) by 10.57 dB (greater than 8-fold). In the control tissue sections, the integrated backscatter from the region of ovarian carcinoma ( $-33.49 \pm 0.86$  dB) was greater ( $p < 0.05$ ) than the normal stromal areas ( $-40.21 \pm 0.61$  dB), approximately 6.72 dB (4-fold), reflecting inherent differences in acoustic character between normal and cancerous ovarian tissue. However, the targeted biotinylated perfluorocarbon emulsion amplified ( $p < 0.05$ ) these inherent differences by approximately 2-fold (3.84 dB; Figure 15). These results clearly demonstrate the ability of site-targeted biotinylated perfluorocarbon emulsion to specifically enhance acoustic detection of ovarian cancer in vitro.

**EXAMPLE 12**

**[0084]** Biotinylated perfluorocarbon emulsion, approximately 250 nm diameter, is targeted to the epithelial capsule of tonsil using monoclonal antibodies specific for cytokeratin, CD-20, and BCL-2 antigens and are acoustically detected using polar, high frequency, high resolution acoustic microscopy. Representative examples of human tonsil are routinely processed by immersion fixation in 10% neutral buffered formalin and embedded in paraffin. Twenty micron sections are prepared for acoustic microscopy; a 5 micron section is used for optical studies. All histologic sections are mounted on acid cleaned glass slides that have been coated with poly-L-lysine. All mounted sections are heated at 55°C for 1 hour in an oven.

**[0085]** Prior to immunostaining, all sections are dewaxed in three changes of Americlear, and dehydrated in successive changes of 95% and 100% ethanol. Endogenous peroxidase activity is blocked only in sections prepared for optical studies by immersion in absolute methanol containing 0.6% (v/v) hydrogen peroxide for 30 minutes. These and all sections for acoustic microscopy are then rehydrated through graded ethanols and distilled water and placed in isotonic PBS (pH 7.4). All sections are incubated with target specific monoclonal antibodies.

**[0086]** Tonsil sections are incubated with a mixture of anti-CD-20, BCL-2, and cytokeratin primary monoclonal antibodies per the recommendations of the vendor for 18 hours at 4°C in moisture chambers. After primary incubation, sections are rinsed in isotonic PBS, then overlain with a polyclonal biotinyl-horse anti-mouse immunoglobulin (VectaStain Elite Kits, Vector Laboratories, Burlingame, CA) for 1 hour at room temperature. After rinsing in PBS, duplicate 30 micron sections are prepared for acoustic microscopy. Sections for light microscopy (5 micron) are incubated with avidin-biotin-peroxidase complex (VectaStain Elite Kit, Vector Lab) for 1 hour at room temperature. Sections are rinsed in phosphate buffer (pH 7.6) and immersed in a solution of 3,3'-diaminobenzidine tetrahydrochloride (Sigma Chemicals, St. Louis, MO; 0.5 mg/ml in phosphate buffer, pH 7.6, containing 0.003% [v/v] hydrogen peroxide) for approximately ten minutes. The chromogenic precipitate are optically enhanced by brief immersion of stained sections in 0.125% (w/v) osmium tetroxide. Sections are then rinsed in tap water, counterstained in Harris' hematoxylin, dehydrated in graded ethanols and Americlear, and mounted in a synthetic mounting medium.

**[0087]** After the second biotinylated antibody is incubated and washed, one slide is incubated in avidin (1.0 mg/-20 cc PBS) using a bath on a rotating table for 30 min. Excess avidin is washed away with isotonic PBS buffer, pH 7.4-7.5 in three 5 minute washes. The prepared slide is incubated with biotinylated perfluorocarbon emulsion for twenty minutes (0.5 cc/-20.0 ml PBS), washed briefly with isotonic PBS 3X for 5 minutes each and rewashed in avidin (1.0 mg/-20 cc) for 15 minutes. Excess avidin is rinsed off with three, 5 minute washes in PBS. The slides are reincubated at above concentrations with biotinylated perfluorocarbon emulsion for 20 minutes. Unbound emulsion is washed away in three changes of PBS (5 minutes each) and the slide is transferred to the acoustic microscope for analysis.

**[0088]** The mounted specimen is immersed into isotonic, phosphate buffered saline at room temperature for ultrasonic insonification. A custom designed acoustic microscope is used to collect ultrasonic data. The microscope consists of a 50 MHz broadband, focused, piezoelectric delay-line transducer (1/4 inch diameter, 1/2 inch focal length, 62 micron beam diameter, Model V390, Panametrics Co., Waltham, MA) operated in the pulse-echo mode. A Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) is used to digitize 35 degree polar backscattered radio frequency (rf) data at 500 megasamples per second with 8 bit resolution. A variable gain system is used to increase the effective dynamic range of this digitizer. Radio frequency data are collected from the entire specimen and a peak detected image is created of the section and compared with the immunostained tissue image.

**[0089]** Immunostained tissue is examined and imaged using a Nikon Optiphot-2 microscope with a Javlin Chromachip 11 camera attachment. Images are routed through a Panasonic digital mixer model WJ-AVE5 to Panasonic SVHS video recorders, models AG- 1960 or AG 1970 and displayed upon an Sony Trinitron monitor. Images are captured using NuVista software (Truevision, Inc., Indianapolis, IN 46256) executing on a Macintosh LCIII microcomputer.

**[0090]** Figure 16 compares tonsil acoustically imaged as a radio frequency peak detected scan at 100 micron lateral step resolution (a) with an optically imaged section immunostained with horseradish peroxidase (b). The epithelial capsule targeted by a mixture of anti-cytokeratin antibodies is distinctly stained with horseradish peroxidase and homologous regions in the acoustic image are "brightened" by the targeted biotinylated acoustic contrast. In Figure 17 the radio frequency peak detected acoustic image at 100 micron step resolution (a) is enhanced

to 50 micron lateral step resolution. The targeted biotinylated perfluorocarbon contrast is clearly seen acoustically enhancing the epithelial rim of the tonsil, analogous to the optical immunostained image. This example clearly demonstrates the fidelity of biotinylated perfluorocarbon contrast targeting for enhanced acoustic contrast of tissues, such as lymph nodes.

### EXAMPLE 13

**[0091]** A Method to Chemically Couple Antibody to Perfluorocarbon Emulsion Particle.

**[0092]** *Preparation of emulsion:* The perfluorocarbon nanoparticle contrast agent is produced by incorporating 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N-4-(pmaleimidophenyl)butyramide (MPB-PE) into the outer lipid monolayer of the emulsion. The emulsion is comprised of perfluorodichlorooctane, safflower oil, a surfactant co-mixture and glycerin. The surfactant co-mixture includes lecithin, cholesterol and MPB-PE which is dissolved in chloroform. The chloroform-lipid mixture is evaporated under reduced pressure, dried in a 50°C vacuum oven overnight and dispersed into water by sonication. The suspension is transferred into a blender cup with perfluorodichlorooctane, *safflower oil* and distilled, deionized water and emulsified for 30 to 60 seconds. The pre-emulsified mixture is transferred to a microemulsifier and continuously processed at 10,000 PSI for three minutes. The completed emulsion is vialled, blanketed with nitrogen and sealed with stopper crimp seal until use. A negative control emulsion is prepared identically, except a nonderivatized phosphatidylethanolamine is substituted into the surfactant co-mixture. Particle sizes are determined in triplicate at 30°C with a laser light scatter submicron particle size analyzer.

**[0093]** *Preparation and Isolation of Tissue Factor F(ab) 'Fragments:* F(ab)' fragments of a specific antibody are generated and isolated using an immunopure F(ab)' preparation kit (Pierce, Rockford, IL). Typically, IgG antibody is dialyzed into 20mM phosphate/10mM EDTA buffer (pH 7.0), concentrated to 20 mg/ml and digested by immobilized papain. Solubilized F9(ab)' are purified from Fc fragments and undigested IgG protein using a protein A column. F(ab)' fragments are



purified from excess cysteine using a G25-15- column and deoxygenated phosphate buffer (pH 6.7). Fraction identity is confirmed by routine SDS-PAGE procedures.

**[0094]**      *Conjugation of Tissue Factor F(ab)' With MPB-PE Derivatized Emulsion:* F(ab)' fractions are pooled and combined with the MPB-PE derivatized emulsion (0.01 to 5.0 mg F(ab)'/ml of emulsion, preferably 1 to 2 mg F(ab)'/ml of emulsion). The mixture is adjusted to pH 6.7, sealed under nitrogen and allowed to react overnight at ambient temperatures with gentle, continuous mixing. The mixture may be subsequently dialyzed with a 300,000 MWCO Spectra/Por DispoDialyzer (Laguna Hills, CA) against 10mM phosphate buffer (PH 7.2) to remove unconjugated F(ab)' fragments. The final emulsion is viald under nitrogen and stored at 4°C until use.

#### EXAMPLE 14

**[0095]**      *In vitro Targeting of Fibrin-Rich Plasma Thrombi Using A Fibrin-Targeted, One-Step Acoustic Contrast System.*

**[0096]**      Whole blood was obtained fresh and anticoagulated (9:1, v/v) with sterile sodium citrate. In a series of trials, plasma clots (9) were produced by combining plasma and 100 mM calcium chloride (3: 1, v/v) with 5 units of thrombin (Sigma Chemical Company, St. Louis, MO) in a plastic tube overlying nitrocellulose membranes. The plasma was allowed to coagulate slowly at room temperature.

**[0097]**      Plasma clots were incubated with anti-fibrin (fab) conjugated or non-conjugated control emulsion contrast using antifibrin monoclonal antibodies (NIB-5F3 or NIB- 1 H 10) (Tymkewycz et al. 1992; Tymkewycz et al. 1993). Half of the clots (5) were incubated individually with 150,ug biotinylated antifibrin monoclonal antibody in 10 ml PBS with 1% bovine serum albumin, (crystallized, Sigma Chemical Company, St. Louis, MO) for two hours; the remaining clots (4) were maintained in PB S with 1 % bovine serum albumin. Bovine serum albumin was added during antibody incubations to minimize nonspecific protein binding to the polystyrene petri dish walls. The anti-fibrin targeted emulsion was incubated with clots (0.2 ml) for 30 minutes. Control clots were treated similarly with a nontargeted control perfluorocarbon emulsion (0.2 ml) for 30 minutes. The plasma clots on

nitrocellulose were insonified using an acoustic microscope to assess the change in ultrasonic backscattered power attributable to the control and targeted emulsions.

**[0098]** The microscope consisted of a 50 MHZ broadband, focused, piezoelectric delay-line transduce (1/4 inch diameter, 1/2 inch focal length, Model V390, Panametrics Co., Waltham, NIA) operated in the pulse-echo mode. A Tektronix DSA 601 digitizing oscilloscope (Beaverton, OR) was used to digitize backscattered radiofrequency data at 500 megasamples per second with 8-bit resolution. Radiofrequency data collected from each site was averaged 32 times. Averaged radiofrequency data were acquired from approximately 400 independent sites with 50 micron lateral step resolution. The radiofrequency data are stored in a low resolution raster scan format and analyzed with custom software. Segments of the radiofrequency lines, 500 nsec in duration and encompassing surface reflection are gated for analysis. The gated data are multiplied by a Hamming window and their power spectra determined by fast-Fourier transformation.

**[0099]** The power spectra from each specimen was referenced to the power spectrum backscattered from a near-perfect steel plate reflector to compute the apparent frequency dependent backscatter transfer function. The backscatter transfer function for the acoustic reflectivity of the smooth cells,  $B(f)$ , was expressed in decibels relative to the power reflected from the steel plate:

$$B(f)^2 = 10 \log [V(f)^2_{\text{tissue}}] / [V(f)^2_{\text{steel plate}}]$$

where  $V(f)^2_{\text{tissue}}$  is the power at selected frequency of the gated rf backscattered from the cells and  $V(f)^2_{\text{steel plate}}$  is the power at the same frequency of the gated rf backscattered from the steel plate. Integrated backscatter (IB) was computed from the average of the frequency-dependent backscatter transfer function over the useful bandwidth of the transducer (Figure 18).

#### EXAMPLE 15

**[00100]** Scanning Electron Micrographs Identification of Tissue Factor Epitopes On Surfaces of Cultured Endothelial Cells Using the One-Step, Tissue Factor Targeted Nanoparticle Emulsion.

**[00101]** Porcine aorta smooth muscle cells known to express tissue factor in abundance on their cell surface were cultured onto 12 mm round, glass cover slips and grown for 48 hours at 37°C under 5% CO<sub>2</sub>. Four treatment groups were prepared: 1) cells alone (untreated control); 2) cells exposed to a nontargeted control

emulsion (negative control); 3) cells exposed to anti-tissue factor antibody before treatment with tissue-factor targeted emulsion (blocked control); and 4) cells treated with tissue-factor targeted emulsion. All smooth muscle cell cultures were fixed in 2.5% glutaraldehyde in PBS for one hour, rinsed briefly in PBS and post-fixed in 2% osmium tetroxide for 1 hour. The fixed and washed cells were dehydrated in ascending concentrations of ethyl alcohol, 10 minutes/step: 50%, 70%, 90% and 3 x 100% followed by three 15 -minute washes in hexamethyldisilazane (Electron Microscopy Sciences, Forth Worth, PA). The glass slips supporting the fixed cells were placed on stubs and imaged with a scanning electron microscope.

**[00102]** Only trace nonspecific binding was appreciated in cell cultures treated with no (Fig. 19A) or untargeted control emulsion (Fig. 19B). Tissue factor targeted specifically bound to the aortic smooth muscle cells (Fig. 19C) and this binding could be specifically blocked by anti-tissue factor antiserum (Fig. 19D). These results demonstrate the ligand specificity of the targeted perfluorocarbon nonospheres.

#### EXAMPLE 16

**[00103]** Local Delivery of One-Step Tissue Factor Targeted Nanoparticle Emulsion In Balloon-Injured Arteries In Pigs:

**[00104]** Briefly, pigs weighing approximately 20 kg were anesthetized with telazol cocktail (1 ml/23 kg IM) followed by 1-2% isoflurane in oxygen. Both common carotids were exposed aseptically; and the distal portion of each, proximal to the bifurcation of the internal carotid artery, was cannulated and connected to a Touhy adapter. The right femoral artery was exposed, and an 8 French sheath was inserted through which a guide catheter and 8 mm balloon catheter was directed to each carotid using an exchange wire. The balloon was inflated five times with sequentially increased pressure up to 6 atmospheres for 30 seconds with one minute between inflations. Prior to the fifth balloon deflation, heparinized (10 units/ml) saline was infused through the distal Touhy access. As the catheter was slowly deflated for the final time and withdrawn from the common carotid, the residual blood within was flushed from the artery. When the catheter was free of the carotid artery, a proximal snare was placed, isolating the carotid segment containing buffer. The anti-tissue factor targeted nanoparticulate emulsion (0.5 ml) was infused into the right carotid

artery and allowed to incubate for 2 hours. After 2 hours the formulation was removed, and the artery was washed with 0.9% NaCl. The left carotid artery was exposed to an nonspecific f(ab) conjugated nanoparticulate emulsion. Vessels were ultrasonically imaged on each side following the formulation washout (Figure 20).

**[00105]** Following angioplasty, but prior to application of contrast agent, carotid arterial walls exhibited no evidence of augmented acoustic reflectivity secondary to overstretch injury alone. Exposure to the control nonspecific emulsion did not alter the acoustic character of the arterial wall. However, the tissue factor targeted emulsion agent clearly penetrated the vascular tissue and localized to the media where increased tissue factor expression was inducted, resulting in substantial augmentation of acoustic contrast. Acoustic enhancement was not circumferentially distributed, but was concentrated over one-half to two-thirds of the vessel wall where non-circumferential force was applied by the angioplasty balloon.

**[00106]** *Immunohistochemistry:* Vessel segments were fixed in molecular biology fixative (Streck Laboratories, Omaha, NE), embedded in paraffin, cut at 5,µm, and mounted on gelatin-coated slides. A slide from each vessel was stained with Verhoeff's van Gieson elastin stain to verify rupture of the internal elastic lamina as an index of deep vascular injury. For immunohistochemistry, the tissue sections were deparaffinized with xylene, hydrated with a descending ethanol series, and rinsed with PBS. Endogenous peroxidase activity was blocked by incubation for 30 minutes with hydrogen peroxide in methanol. Nonspecific binding of antibody was blocked by incubation for 45 minutes in PBS containing 1% bovine serum albumin, 0.3% triton X-100, 10% normal goat serum, and 1% normal pig serum. The sections were incubated overnight at 4°C with the rabbit anti-pig tissue factor antibody diluted 1: 100 in the blocking solution. After rinsing the slides with PBS, the sections were incubated for 45 minutes at room temperature with biotinylated goat anti-rabbit IgG (Vector Laboratories) diluted 1:200 with blocking buffer. The sections were stained with AEC (HistostainPlus, Zymed) and counterstained with hematoxylin.

**[00107]** Immunohistochemistry revealed a pattern of localization of tissue factor in balloon-injured vessels in the media (arrows) at the site of acoustic contrast enhancement, as well as on the intimal surface associated with inflammatory cells (Fig. 21A; L=lumen). In contrast, control uninjured vessels exhibited staining primarily in the adventitia and not in the tunica media (Fig. 21B). No obvious dissections of the arterial walls were observed microscopically, although some mild

fenestration of the internal elastic lamina was noted. These results demonstrate the presence of tissue factor only in the media of balloon overstretched arteries. The vessels were acoustically enhanced by the tissue factor targeted, one-step nanoparticulate emulsion, but not by the nontargeted control nanoparticulate emulsion.

**[00108]** In view of the above, it will be seen that the several objects of the invention are achieved and other advantageous results attained.

**[00109]** As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description and shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
21